

Parametric excitation of plasma waves by gravitational radiation

Gert Brodin* and Mattias Marklund†

Department of Plasma Physics, Umeå University, S-901 87 Umeå, Sweden

We consider the parametric excitation of a Langmuir wave and an electromagnetic wave by gravitational radiation, in a thin plasma on a Minkowski background. We calculate the coupling coefficients starting from a kinetic description, and the growth rate of the instability is found. The Manley–Rowe relations are fulfilled only in the limit of a cold plasma. As a consequence, it is generally difficult to view the process quantum mechanically, i.e. as the decay of a graviton into a photon and a plasmon. Finally we discuss the relevance of our investigation to realistic physical situations.

04.30.Nk, 52.35.Mw, 95.30.Sf

The state of matter in regions where general relativistic treatments are desirable is often the plasma state. Nevertheless, plasma physics and general relativity are quite distinct areas of physics, and accordingly there are comparatively few papers using a general relativistic framework that include the electromagnetic forces in their treatments. However, there are a few exceptions to this rule, see for example Refs. 1–5, where the plasma dynamics in a strong gravitational field is considered [1,2], relativistic transport equations for a plasma are derived [3], a general relativistic version of the Kelvin–Helmholtz theorem is derived [4] and photon acceleration by gravitational radiation is considered [5]. Omission of the electromagnetic effects for a plasma subject to gravitational forces is possible because gravity alone do not separate the charges, and in many cases the plasma can be treated as a neutral fluid, in spite of its electromagnetic properties.

In this letter we will consider a simple model problem, that has two interesting properties: Firstly, the process of investigation requires a general relativistic description of the plasma dynamics to occur, and secondly we demonstrate the possibility of charge separation induced by the gravitational effects. We start from a monochromatic gravitational wave propagating through a thin plasma superimposed on a flat background metric, and consider the parametric excitation of a plasma wave and an electromagnetic wave. The plasma wave - that has an increasing amplitude due to the above mentioned mechanism - undergoes charge density oscillations. Thus the parametric excitation process - which of course has no Newtonian analog - shows that although no *direct* charge separation can be caused by gravitational forces, *indirectly* charge density perturbations may grow due to the gravitational and electromagnetic interaction. Theoretical aspects of our problem, i.e. satisfaction of the Manley–Rowe relations and the possibility of a quantum interpretation of the interaction, and the relevance of our model problem to realistic physical situations will be discussed in the end of this letter.

We will refer to tensorial objects in terms of their components in a coordinate basis. Note that as observers the natural way to measure quantities is to project them onto a Lorentz frame $\{\mathbf{e}_{\hat{a}}, \hat{a} = 0, \dots, 3\}$, i.e., a frame in which the metric components become those of the Minkowski metric. The frame components of the electric and magnetic fields $E^{\hat{a}}$ and $B^{\hat{a}}$ are defined by $F^{\hat{0}\hat{a}} = E^{\hat{a}}$, $F^{\hat{a}\hat{b}} = \delta_{\hat{c}\hat{d}} \epsilon^{\hat{a}\hat{b}\hat{c}} B^{\hat{d}}$, where $\epsilon^{\hat{a}\hat{b}\hat{c}}$ are the components of the totally skew tensor. Expanding $\mathbf{E} = E^{\hat{a}} \mathbf{e}_{\hat{a}}$ and $\mathbf{B} = B^{\hat{a}} \mathbf{e}_{\hat{a}}$ to first order in the metric's coordinate components, we get that the electric field components in the coordinate basis are the same as in the Lorentz frame. Furthermore, the magnetic field components deviates only to first order in the metric components from the Lorentz tetrad components, and it is therefore meaningful to use the coordinate components in place of the tetrad components. This would of course not be the case in a strong gravitational field such as, say, the Kerr geometry.

The Vlasov equation for the distribution $f = f(x^\mu, p^a)$ (where $\mu, \nu, \dots = t, x, y, z$ and $a, b, \dots = x, y, z$) reads [6]

$$\frac{\partial f}{\partial t} + \frac{p^a}{p^t} \frac{\partial f}{\partial x^a} + \left(\frac{q}{p^t} g_{\mu\nu} F^{\mu\nu} p^\nu - \mathcal{G}^a \right) \frac{\partial f}{\partial p^a} = 0, \quad (1)$$

where $\mathcal{G}^a \equiv \Gamma_{\mu\nu}^a p^\mu p^\nu / p^t$, and this is coupled to Maxwell's equations

$$F^{\mu\nu}_{;\nu} = \mu_0 j^\mu \equiv \sum_{\text{p.s.}} \mu_0 q \int f(x^\nu, p^a) p^\mu \frac{|g|^{1/2}}{|p_t|} d^3 p, \quad (2a)$$

*E-mail address: gert.brodin@physics.umu.se

†E-mail address: mattias.marklund@physics.umu.se

$$F_{[\mu\nu,\sigma]} = 0 , \quad (2b)$$

where p.s. stands for particle species. Here we have introduced the invariant measure $(|g|^{1/2}/|p_t|)d^3p$ on the surface in momentum space where $g_{\mu\nu}p^\mu p^\nu = -m^2$, m being the mass of the particle in question, and we use g to denote the determinant of the metric. The gravitational field, represented by the Christoffel symbols $\Gamma_{\nu\sigma}^\mu$ in the Vlasov equation, is assumed to be generated by some outside source. The plasma itself is assumed to generate a much weaker gravitational field.

Next we assume the presence of a small amplitude gravitational wave, and we use the transverse traceless gauge. The line element then takes the form [7]

$$ds^2 = -dt^2 + (1 + h_+) dx^2 + (1 - h_+) dy^2 + 2h_\times dx dy + dz^2 , \quad (3)$$

where $h_+ = h_+(u)$, $h_\times = h_\times(u)$, with $u \equiv z - t$, and $|h_+|, |h_\times| \ll 1$. The mass of a particle with momentum p^μ is defined as $m \equiv |g_{\mu\nu}p^\mu p^\nu|^{1/2}$. From this, p^t may be expressed in terms of m and the 3-momentum p^a :

$$p^t = \{m^2 + \delta_{ab}p^a p^b + h_+ [(p^x)^2 - (p^y)^2] + 2h_\times p^x p^y\}^{1/2} . \quad (4)$$

To first order in h_+ , $h_\times$, the Christoffel symbols read

$$-\Gamma_{xx}^t = \Gamma_{yy}^t = -\Gamma_{tx}^x = \Gamma_{xz}^x = \Gamma_{ty}^y = -\Gamma_{yz}^y = -\Gamma_{xx}^z = \Gamma_{yy}^z = \frac{1}{2}\dot{h}_+ , \quad (5a)$$

$$-\Gamma_{xy}^t = -\Gamma_{ty}^x = \Gamma_{yz}^x = -\Gamma_{tx}^y = \Gamma_{xz}^y = -\Gamma_{xy}^z = \frac{1}{2}\dot{h}_\times , \quad (5b)$$

where $\dot{h} \equiv dh/du$. Using the Christoffel symbols (5) of the gravitational wave, the Vlasov equation (1) for the unperturbed ($F^{\mu\nu} = 0$) distribution $f = f(t, x^a, p^b)$ becomes

$$\frac{\partial f}{\partial t} + \frac{p^a}{p^t} \frac{\partial f}{\partial x^a} - \mathcal{G}^a \frac{\partial f}{\partial p^a} = 0 , \quad (6)$$

where

$$\begin{aligned} \mathcal{G}^x &\equiv \left(-1 + \frac{p^z}{p^t}\right) (\dot{h}_+ p^x + \dot{h}_\times p^y) , \\ \mathcal{G}^y &\equiv \left(-1 + \frac{p^z}{p^t}\right) (\dot{h}_\times p^x - \dot{h}_+ p^y) , \\ \mathcal{G}^z &\equiv \frac{1}{2p^t} (\gamma_{AB} p^A p^B \dot{h}_+ - 2p^x p^y \dot{h}_\times) , \end{aligned}$$

and $(\gamma_{AB}) = \text{diag}(-1, 1)$, $A, B = x, y$. Expanding p^t according to $p^t \approx p_{(0)}^t - \mathcal{F}$, where $\mathcal{F} \equiv (h_+ \gamma_{AB} p^A p^B - 2h_\times p^x p^y) / (2p_{(0)}^t)$, and $p_{(0)}^t = (m^2 + \delta_{ab} p^a p^b)^{1/2}$, the Vlasov equation becomes

$$\frac{\partial f}{\partial t} + \left(1 + \frac{\mathcal{F}}{p_{(0)}^t}\right) \frac{p^a}{p_{(0)}^t} \frac{\partial f}{\partial x^a} - \mathcal{G}^a \frac{\partial f}{\partial p^a} = 0 , \quad (7)$$

For simplicity we let the gravitational wave be monochromatic:

$$h = \tilde{h} \exp[i(k_0 z - \omega_0 t)] + \text{c.c.} , \quad (8)$$

with the dispersion relation $k_0 = \omega_0$. We divide f according to $f = f_{\text{SJ}}(p^a) + f_g(t, z, p^a)$ where the equilibrium distribution f_{SJ} is taken to be the Sygne-Jüttner distribution [8], and $f_g = \tilde{f}_g \exp[i(k_0 z - \omega_0 t)]$ is the perturbation induced by the gravitational wave - which is assumed to fulfill $\tilde{f}_g \ll f_{\text{SJ}}$. From Eq. (7) we find

$$\tilde{f}_g = - \frac{\mathcal{G}^a (\partial f_{\text{SJ}} / \partial p^a)}{i(\omega_0 - k_0 p^z / p_{(0)}^t)} \quad (9)$$

to first order in the amplitude.

Next we assume the presence of electromagnetic perturbations with frequency and wavenumber $(\omega_1, \mathbf{k}_1)$ as well as electrostatic perturbations $(\omega_2, \mathbf{k}_2)$. The frequencies and wavenumbers are taken to satisfy the resonance conditions

$\omega_0 = \omega_1 + \omega_2$ and $\mathbf{k}_0 = \mathbf{k}_1 + \mathbf{k}_2$. The vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ span a plane, and we chose the y -axis to be perpendicular to this plane, i.e. $\mathbf{k}_1 = k_1^x \hat{\mathbf{x}} + k_1^z \hat{\mathbf{z}}$ and $\mathbf{k}_2 = k_2^x \hat{\mathbf{x}} + k_2^z \hat{\mathbf{z}}$. Including the electromagnetic field, expanding Eq. (7) to first order in h and writing all terms proportional to h on the right hand side Eq. (7) becomes

$$\begin{aligned} & \frac{\partial f_{\text{ng}}}{\partial t} + \frac{p^a}{p_{(0)}^t} \frac{\partial f_{\text{ng}}}{\partial x^a} + q \left(-F^{at} + \frac{1}{p_{(0)}^t} \delta_{bc} F^{ac} p^b \right) \frac{\partial f_{\text{ng}}}{\partial p^a} \\ &= -\frac{\mathcal{F}}{(p_{(0)}^t)^2} p^a \frac{\partial f_{\text{ng}}}{\partial x^a} + \mathcal{G}^a \frac{\partial f_{\text{ng}}}{\partial p^a} - q \left(-F^{at} + \frac{1}{p_{(0)}^t} \delta_{bc} F^{ac} p^b \right) \frac{\partial f_{\text{g}}}{\partial p^a} \\ & - \left\{ \frac{q}{p_{(0)}^t} [-h_+ \gamma_{BC} F^{aC} p^B + h_{\times} (F^{ax} p^y + F^{ay} p^x)] \right\} \frac{\partial f_{\text{SJ}}}{\partial p^a} \end{aligned} \quad (10)$$

where we have introduced f_{ng} defined by $f_{\text{ng}} = f - f_{\text{g}}$. Furthermore, we need two components of Faraday's law. Expanding Eq. (2b) to first order in the amplitude h , we obtain

$$\partial_t F^{xy} - \partial_y F^{tx} + \partial_x F^{ty} = h_{\times} (-\partial_x F^{tx} + \partial_y F^{ty}) + h_+ (\partial_y F^{tx} + \partial_x F^{ty}) , \quad (11a)$$

$$\begin{aligned} \partial_t F^{yz} - \partial_z F^{ty} + \partial_y F^{tz} &= h_{\times} (\partial_z F^{tx} - \partial_t F^{xz}) + h_+ (\partial_z F^{ty} + \partial_t F^{yz}) \\ &+ \dot{h}_{\times} (F^{tx} + F^{xz}) - \dot{h}_+ (F^{ty} + F^{yz}) , \end{aligned} \quad (11b)$$

In what follows, we will put $h_+ = 0$, i.e., we choose a specific polarization of the gravitational wave. It turns out that such a gravitational wave does not couple to a linearly polarized electromagnetic wave with magnetic field in the y -direction. Thus we assume the electromagnetic wave to have the opposite polarization, i.e. F^{yt}, F^{xz} and F^{yz} are the only electromagnetic components to be different from zero in the linear approximation. Similarly F^{zt} and F^{xt} are the only nonzero electrostatic components of the field tensor. We then divide f_{ng} according to $f_{\text{ng}} = f_{\text{SJ}}(p^a) + \widetilde{f_{\text{em}}}(t, p^a) \exp[i(k_1^x x + k_1^z z - \omega_1 t)] + \widetilde{f_{\text{es}}}(t, p^a) \exp[i(k_2^x x + k_2^z z - \omega_2 t)]$ where the time dependence of the amplitudes, which is due to the parametric interaction with the gravitational wave, is assumed to be slow, such that $\partial/\partial t \ll \omega$. In the linear approximation (i.e. no gravitational coupling) the slow time dependence vanishes, and Eq. (10) gives

$$\widetilde{f_{\text{em}}}^{\text{l}} = -\frac{q \widetilde{F}^{yt}}{i\widehat{\omega}_1} \frac{\partial f_{\text{SJ}}}{\partial p^y} , \quad (12a)$$

$$\widetilde{f_{\text{es}}}^{\text{l}} = -\frac{q \widetilde{F}^{at}}{i\widehat{\omega}_2} \frac{\partial f_{\text{SJ}}}{\partial p^a} \quad (12b)$$

for the electromagnetic and electrostatic perturbations respectively. Here we have $\widehat{\omega}_i \equiv \omega_i - \mathbf{k}_i \cdot \mathbf{p}/p_{(0)}^t$, where the scalar product is defined as $\mathbf{A} \cdot \mathbf{B} \equiv \delta_{ab} A^a B^b$. Considering the part of Eq. (10) varying as $\exp[i(k_2^x x + k_2^z z - \omega_2 t)]$ using linear approximations for the factors on the right hand side we obtain $\widetilde{f_{\text{es}}} = \widetilde{f_{\text{es}}}^{\text{l}} + \widetilde{f_{\text{es}}}^{\text{nl}}$. The linear contribution is given by (12b), by replacing ω_2 by $\omega_{2\text{op}} \equiv \omega_2 + i\partial/\partial t$ (since $\partial/\partial t \ll \omega_2$, division by $\omega_{2\text{op}}$ can be calculated by a first order Taylor expansion) and the nonlinear contribution is found to be $\widetilde{f_{\text{es}}}^{\text{nl}} = C_{\text{es}} \widetilde{h}_{\times} \widetilde{F}^{yt^*}$ where

$$\begin{aligned} C_{\text{es}} &= \frac{q}{i\widehat{\omega}_2} \left\{ \frac{p^x p^y \mathbf{k}_1 \cdot \mathbf{p}}{(p_{(0)}^t)^3 \widehat{\omega}_1} \frac{\partial f_{\text{SJ}}}{\partial p^y} + \omega_0 \mathcal{G}_{\text{op}} \left(\frac{1}{\widehat{\omega}_1} \frac{\partial f_{\text{SJ}}}{\partial p^y} \right) \right. \\ & \left. + \left[\left(1 - \frac{1}{p_{(0)}^t} \left(\frac{\mathbf{p} \cdot \mathbf{k}_1}{\omega_1} \right) \right) \frac{\partial}{\partial p^y} + \frac{p^y}{\omega_1 p_{(0)}^t} \left(k_1^x \frac{\partial}{\partial p^x} + k_1^z \frac{\partial}{\partial p^z} \right) \right] \left[\frac{\omega_0}{\widehat{\omega}_0} (\mathcal{G}_{\text{op}} f_{\text{SJ}}) \right] + \frac{k_1^z p^x}{p_{(0)}^t \omega_1} \frac{\partial f_{\text{SJ}}}{\partial p^z} \right\} \end{aligned} \quad (13)$$

and $\mathcal{G}_{\text{op}} \equiv (1 - p^z/p_{(0)}^t)(p^y \partial/\partial p^x + p^x \partial/\partial p^y) + (p^x p^y/p_{(0)}^t) \partial/\partial p^z$. Combining Eqs. (13) and (2a) we then find

$$\varepsilon_L(\omega_{2\text{op}}, k_2) \widetilde{F}^{es} = \frac{\mu_0 q \widetilde{h}_{\times} \widetilde{F}^{yt^*}}{ik_2} \sum_{\text{p.s.}} \int C_{\text{es}} d^3 p. \quad (14)$$

where the longitudinal dielectric permittivity ε_L is given by [9]

$$\varepsilon_L = 1 + \frac{\mu_0 q}{k_2} \sum_{\text{p.s.}} \int \frac{1}{\widehat{\omega}_{2\text{op}}} \frac{\partial f_{\text{SJ}}}{\partial p} d^3 p, \quad (15)$$

$p = [(p^x)^2 + (p^y)^2 + (p^z)^2]^{1/2}$, $k_2 = [(k_2^x)^2 + (k_2^z)^2]^{1/2}$, and $\tilde{F}^{es} = [(\tilde{F}^{xt})^2 + (\tilde{F}^{zt})^2]^{1/2}$. In deriving Eqs. (14) and (15) we have used that the electrostatic mode is longitudinal, i.e. $\tilde{F}^{xt}/\tilde{F}^{zt} = k_2^x/k_2^z$.

Next we turn to the part of (10) varying as $\exp[i(k_1^x x + k_1^z z - \omega_1 t)]$. Again using linear approximations for the right hand side we obtain $\widetilde{f_{\text{em}}} = \widetilde{f_{\text{em}}^l} + \widetilde{f_{\text{em}}^{\text{nl}}}$ where $\widetilde{f_{\text{em}}^l}$ is given by (12a) (naturally by replacing ω_1 with $\omega_{1\text{op}}$), and $\widetilde{f_{\text{em}}^{\text{nl}}}$ is found to be $\widetilde{f_{\text{em}}^{\text{nl}}} = C_{\text{em}} \tilde{h}_{\times} \tilde{F}^{es*}$ where

$$C_{\text{em}} = \frac{q}{i\hat{\omega}_1} \left\{ \omega_0 \mathcal{G}_{\text{op}} \left(\frac{\hat{k}_2^x}{\hat{\omega}_2} \frac{\partial f_{\text{SJ}}}{\partial p^x} + \frac{\hat{k}_2^z}{\hat{\omega}_2} \frac{\partial f_{\text{SJ}}}{\partial p^z} \right) - \left(\hat{k}_2^x \frac{\partial}{\partial p^x} + \hat{k}_2^z \frac{\partial}{\partial p^z} \right) \left[\frac{\omega_0}{\hat{\omega}_0} \mathcal{G}_{\text{op}} f_{\text{SJ}} \right] - \frac{p^x p^y \mathbf{k}_2 \cdot \mathbf{p}}{(p_{(0)}^t)^3} \left(\frac{\hat{k}_2^x}{\hat{\omega}_2} \frac{\partial f_{\text{SJ}}}{\partial p^x} + \frac{\hat{k}_2^z}{\hat{\omega}_2} \frac{\partial f_{\text{SJ}}}{\partial p^z} \right) \right\}. \quad (16)$$

Eliminating linear terms proportional to F^{yz} and F^{xy} with the help of Eqs. (11b) and (11a), Eq. (2a) together with (16) gives

$$\begin{aligned} D(\omega_{1\text{op}}, |\mathbf{k}_1|) \tilde{F}^{yt} &\equiv \left(\varepsilon_T(\omega_{1\text{op}}, |\mathbf{k}_1|) - \frac{\mathbf{k}_1^2 c^2}{\omega_{1\text{op}}^2} \right) \tilde{F}^{yt} \\ &= \left\{ \frac{i\mu_0 q}{\omega_1} \int \left[\frac{p^y}{p_{(0)}^t} \left(C_{\text{em}} + \frac{qp^x p^y}{i(p_{(0)}^t)^2} \left(\frac{k_2^x}{k_2 \hat{\omega}_2} \frac{\partial f_{\text{SJ}}}{\partial p^x} + \frac{k_2^z}{k_2 \hat{\omega}_2} \frac{\partial f_{\text{SJ}}}{\partial p^z} \right) \right) \right] d^3 p + \frac{k_{2x} k_1^2}{k_2 \omega_1^2} \right\} \tilde{h}_{\times} \tilde{F}^{es*} \end{aligned} \quad (17)$$

where the transverse dielectric permittivity ε_T is given by [9]

$$\varepsilon_T = 1 + \frac{\mu_0 q^2}{\omega_1} \sum_{\text{p.s.}} \int \frac{p^y}{\hat{\omega}_1} \frac{\partial f_{\text{SJ}}}{\partial p^y} d^3 p. \quad (18)$$

The operators $\varepsilon_L(\omega_{2\text{op}}, k_2)$ and $D(\omega_{1\text{op}}, |\mathbf{k}_1|)$ can be expanded according to $\varepsilon_L(\omega_{2\text{op}}, k_2) = \varepsilon_L(\omega_2, k_2) + (\partial \varepsilon_L / \partial \omega_2) i \partial / \partial t$ and $D(\omega_{1\text{op}}, |\mathbf{k}_1|) = D(\omega_1, |\mathbf{k}_1|) + (\partial D / \partial \omega_1) i \partial / \partial t$. Assuming that the dispersions relations for the Langmuir wave and the electromagnetic wave are exactly fulfilled, we let $\varepsilon_L(\omega_2, k_2) = D(\omega_1, |\mathbf{k}_1|) = 0$. Equations (14) and (17) are then written

$$\frac{\partial \tilde{F}^{es}}{\partial t} = C_1 \frac{\tilde{h}_{\times} \tilde{F}^{yt*}}{(\partial \varepsilon_L / \partial \omega_2)}, \quad (19a)$$

$$\frac{\partial \tilde{F}^{yt}}{\partial t} = C_2 \frac{\tilde{h}_{\times} \tilde{F}^{es*}}{(\partial D / \partial \omega_1)} \quad (19b)$$

where the (constant) coupling coefficients C_1 and C_2 are trivially found from (14) and (17). As can be seen, generally the expressions for the coupling coefficients are very complicated. In order to get transparent formulas, we present the result for a cold electron plasma with immobile ions constituting a neutralizing background. Letting $f_{\text{SJ}} \rightarrow \delta^3(\mathbf{p})$ the coupling coefficients become $C_1 = C_2 \equiv C = -ik_{2x}\omega_0/k_2\omega_1$. Using these expressions and combining (19a) and (19b) the growth rate γ for the Langmuir wave and the electromagnetic wave is given by

$$\gamma^2 = \left(\frac{k_{2x}\omega_0}{k_2\omega_1} \right)^2 \frac{\omega_1\omega_2}{4} |\tilde{h}_{\times}|^2. \quad (20)$$

Since we have not included dissipation of the wave modes, the threshold value for the instability is zero. However, it is straightforward [10] to calculate the threshold by including appropriate damping mechanisms for the decay products.

The fact that the same coupling coefficient C appears in Eqs. (19a) and (19b) for a cold plasma means that the Manley–Rowe relations [10] are fulfilled for that case. These relations usually follows from an underlying Hamiltonian structure of the governing equations, and assures that each of the decay products takes energy from the pump wave in direct proportion to their respective frequencies. Thus fulfillment of the Manley–Rowe relation means that the parametric process can be interpreted quantum mechanically - i.e. we can think of the interaction as the decay of a graviton with energy $\hbar\omega_0$ into a photon with energy $\hbar\omega_1$ and a plasmon with energy $\hbar\omega_2$. However, we stress that this interpretation is not always possible. When thermal effects are taken into account, the coefficient C_1 in (19a) is generally different from the coefficient C_2 in (19b). Thus we obtain the rather surprising result that the simple

graviton interpretation (i.e. that the gravitational wave is built up of quanta with energy $\hbar\omega_0$) of the gravitational field is not always applicable.

To our knowledge, our set of governing Eqs. (1)–(2), have not been used to consider specific physical processes previously. Probably this is because the system has a very high complexity - it is nonlinear and has seven independent variables - but, on the other hand, simpler models for the plasma cannot deal with thermal effects properly. As a starting point we have addressed a simple "model problem", to reduce the complexity of our governing equations. The main conclusion that can be drawn from our calculations are two: Firstly, wave-wave interactions can lead to energy transfer from gravitational to electromagnetic degrees of freedom, and vice versa. Secondly, charge separation and corresponding electrostatic fields may occur as a result of such interactions. However, the idealizations made in our calculations make it difficult to make specific physical predictions. For example, binary systems can produce monochromatic gravitational radiation, but the amplitude is probably too low to overcome the threshold value caused by plasma inhomogeneities [11], except possibly in the very last collapse phase when the system produces an incoherent broadband gravitational pulse. Similarly, supernova events may produce high enough gravitational wave amplitudes for significant energy transfer to electromagnetic degrees of freedom to occur, but in most cases we expect the gravitational radiation to be very incoherent and broadband (there are calculations predicting highly monochromatic emissions of gravitational waves for certain types of neutron star formation [12]). Some estimates of the effects due to scattering of gravitational waves in supernova has been presented by Bingham *et al.* [13]. One may also consider the formation of black holes as a source of gravitational radiation, especially the non-axisymmetric collapse, but - again - this process gives a pulse rather than a monochromatic wave. (See the review by Thorne [14] for characteristic values of the amplitudes, frequencies, and duration times for the processes discussed above.)

The parametric process considered in this letter can be thought of as a gravitational analog of Raman scattering, where the electromagnetic pump wave has been replaced by a gravitational wave. Similarly, we can imagine that a gravitational analog of Compton scattering may occur. Such a process requires a kinetic treatment, and could be described within our formalism without very much extra difficulty. Gravitational waves may also be subject to four wave processes such as modulational instabilities. Furthermore, a gravitational radiation background similar to the microwave background is assumed to exist as a relic from big bang [15]. In the early universe the intensity of such radiation could be very high, and - if the magnitude of the gravitational waves are sufficient - processes such as those mentioned above could be of importance for the structure formation in the universe. It should be stated though that estimates of the amplitude of such primordial gravitational waves are very difficult to make, much depending on the insufficient data concerning the early universe. Also, much of the simplifying assumption made in this letter must be relaxed before such issues can be studied seriously.

-
- [1] K. Holcomb and T. Tajima (1989) *Phys. Rev. D* **40** 3809.
 - [2] J. Daniel and T. Tajima (1997) *Phys. Rev. D* **55** 5193.
 - [3] A. Georgiou (1995) *Class. Quantum Grav.* **12** 1491.
 - [4] K. Elsässer and S. Popel (1997) *Phys. Plasmas* **4** 2348.
 - [5] J. T. Mendonça, P. K. Shukla and R. Bingham (1998) Photon Acceleration by Gravitational Waves, submitted to *Phys. Lett.*
 - [6] J. Ehlers (1971) General Relativity and Kinetic Theory, in *General Relativity and Cosmology*, Proceedings of the International School of Physics Enrico Fermi, Course XLVII, ed. R. K. Sachs (Academic Press).
 - [7] C. W. Misner, K. S. Thorne, and J. A. Wheeler (1973) *Gravitation* (W. H. Freeman and Company).
 - [8] R. L. Liboff (1998) *Kinetic Theory* (John Wiley & Sons Inc.)
 - [9] G. Brodin and L. Stenflo (1989) *J. Plasma Physics* **42** 187.
 - [10] J. Weiland and H. Wilhelmsson (1977) *Coherent Non-Linear Interaction of Waves in Plasmas* (Pergamon Press).
 - [11] K. Nishikawa and C. S. Liu (1976) in *Advances in Plasma Physics* 6, eds. A. Simon and W. B. Thompson (Wiley, New York, London, Sydney, Toronto).
 - [12] J. R. Ipser and R. A. Managan (1984) *Astrophysical Journal* **282** 287.
 - [13] R. Bingham, R. A. Cairns, J. M. Dawson, R. O. Dendy, C. N. Lashmore-Davies, J. T. Mendonça, P. K. Shukla, L. O. Silva and L. Stenflo (1998) *Phys. Scripta* **T75** 61.
 - [14] K. S. Thorne (1987) in *Three hundred years of gravitation*, eds. S. W. Hawking and W. Israel (Cambridge University Press).
 - [15] P. J. E. Peebles (1993) *Principles of Physical Cosmology* (Princeton University Press).